

**POISON**

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

FARMALINX

**OxaFert**

**Herbicide and Fertiliser**

**ACTIVE CONSTITUENT: 10 g/kg OXADIAZON**

**SOLVENT: 10.5 g/kg POLYBUTENE**

**GROUP G HERBICIDE**

*Combination herbicide and fertiliser for pre-emergent control of summer grass & crab grass and winter grass in turf, feeding of turf and for the control of certain weeds in woody ornamental shrubs in nurseries.*

**GUARANTEED FERTILISER ANALYSIS**

12.0%	TOTAL NITROGEN (N)
3.0%	As ammoniacal nitrogen (N)
9.0%	As nitrate nitrogen (N)
5.0%	PHOSPHORUS (P) available phosphoric acid
14.0%	POTASSIUM (K) soluble potash
8.0%	SULPHUR (S) as elemental
1.5%	MAGNESIUM (Mg)
2.0%	CALCIUM (Ca)
0.02%	BORON (B)
0.1%	ZINC (Zn)
0.02%	IRON (Fe)
0.02%	MOLYBDENUM (Mo)

**farmalinx** *Partners in Agriculture*

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844 W: www.farmalinx.com

NET CONTENTS

**20 kg**

**farmalinx**  
*Partners in Agriculture*

**OXAFERT**

**DIRECTIONS FOR USE**

**RESTRAINTS:**

**DO NOT** apply granules when the turf foliage is wet.

**DO NOT** use on putting greens, low cut tees or bowling greens.

**DO NOT** apply to newly seeded turf areas.

Situation	Target	Turf Species	Rate	Critical Comments
Warm season turfgrass areas, including fairways, golf tees, municipal sports fields, industrial and commercial lawns	Pre emergent control of Summer Grass ( <i>Digitaria ciliaris</i> ), Crowsfoot Grass ( <i>Elcusine indica</i> )	Kikuyu ( <i>Pennisetum clandestinum</i> ), Buffalo Grass ( <i>Stenotaphrum secundatum</i> ), Common Couch ( <i>Cynodon dactylon</i> ), <i>C. transvaalensis</i> ) Grass (Not Santa Ana) Tif Hybrid Couch Grass, Saltwater couch ( <i>Paspalum vaginatum</i> ), Queensland Blue Couch ( <i>Digitaria didactyla</i> ), Bahia grass ( <i>Paspalum notatum</i> ), Zoysia grass ( <i>Zoysia matrella</i> )	300 kg/ha (3 kg/100 m <sup>2</sup> )	The product is ready to apply. No mixing or dilution is required. Apply product evenly to dry foliage and bare areas using a granular spreader able to give even coverage. Application must be made before weeds have begun to germinate. Established weeds will not be controlled. After application water in with at least 10mm of sprinkler irrigation. Control of germinating weed seedlings can be expected for up to 10 weeks after application depending upon temperature, rainfall, soil type and other factors. If longer control is required a repeat application should be made 7-10 weeks after initial application. <b>DO NOT</b> mow treated area until product has been watered in and turf foliage has dried. Avoid application when daytime temperatures are consistently above 32°C. This product is not recommended for use on putting greens, low cut tees or bowling greens. <b>DO NOT</b> apply on turf containing Carpet Grass ( <i>Axonopus</i> spp.), Centipede Grass ( <i>Eremochloa ophiuroides</i> ) and Dichondra unless some damage is acceptable.
	Pre emergent Control of Winter Grass ( <i>Poa annua</i> ) Creeping Oxalis ( <i>Oxalis comiculala</i> )		400 kg/ha (4 kg/100 m <sup>2</sup> )	
	Turf nutrition - provides slow release nitrogen for 8 weeks feeding		300-400 kg/ha (3-4 kg/100m <sup>2</sup> )	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**GENERAL INSTRUCTIONS**

Farmalinx OxaFert Herbicide and Fertiliser is a convenient ready to use formulation for the control of certain annual grass weeds in turf and also for fertilisation of turf where nitrogen, potassium and sulphur are required. The product should only be used when both functions are required.

**APPLICATION**

For a granular spreader which can be calibrated for even output and capable of giving even application. Use a double application technique: i.e. half rate width, to achieve a full rate application. Apply under calm conditions, thereby reducing drift to non-target areas and optimizing particle distribution for good weed control.

Ensure turf area is free of leaves and other debris, mowing if necessary so that the granules reach the soil surface. Delay reseeding of Ryegrass or Bentgrass into treated areas for 4-5 months (Ryegrass 4 months & Bentgrass 5 months).

**DO NOT** over apply, especially under the branch spread of trees and shrubs.

Avoid contact with flowers and shrubs.

Suggested Spreader Settings				
The suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product.				
Spreader	Ground Speed	Width of coverage	Spreader Settings	
			300 kg/ha	400 kg/ha
Scotts AccuPro 2000, SR-2000 (cone 4)	4.8 km/hr	4.0 m	0 1/2	Q
Scotts R-8, R-8A (cone 4)	4.8 km/hr	4.0 m	0	P 1/2
Spyker 288 Super	4.8 km/hr	3.0 m	3.9	4.2
Vicon (03 Series - Standard Spout)	7.3 km/hr	11.0 m	37	44

**HERBICIDE RESISTANCE WARNING**

Farmalinx OxaFert Herbicide and Fertiliser is a member of the Oxadiazole Group of herbicide.

**GROUP G HERBICIDE**

The herbicide is an inhibitor of protoporphyrinogen oxidase. For weed resistance management, the product is a Group G Herbicide. Some naturally occurring weed biotypes resistant to the herbicide may exist through normal genetic variability in weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other oxadiazole group herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd does not accept liability for any losses that may result from failure of this product to control resistant weeds.

APVMA Approval No.: 70376/63316

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Dangerous to Fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply product if rapid storm run-off from treated areas into water bodies is likely to occur within several days of application.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Do not store with seed, feed, foodstuffs, insecticides, fungicides or other fertilisers. Shake bag empty into applicator. Puncture and shred empty container and bury in a local authority landfill. If no landfill is available bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT dispose of damaged material on site. DO NOT re-use this container for any purpose.

**SPILLS**

Sweep up spilled materials and return to original container. Wash contaminated equipment taking care the washings DO NOT enter sewers, drains and other waterways.

**SAFETY DIRECTIONS**

Avoid contact with eyes. Do not inhale dust. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX OxaFert Herbicide and Fertiliser is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

**NOTICE TO BUYER**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.



20 kg

Batch No.

Date of Manufacture

**Farmalinx**  
 Partners in Agriculture

**OXAFERT**