LOOK WHO'S BACK





THE MOST

TRUSTED

BROADLEAF

HERBICIDES



COLIN CAMPBELL (CHEMICALS) PTY LTD • (02) 9725 2544 HORTICULTURE • POST HARVEST • TURF

80 Years in 2020 • Est. 1940 • campbellchemicals.com.au 100% Australian owned & operated.





At a Glance



Product Type:

Resistance group:

Pack Size:

Herbicide

Group 4 Herbicide

20L (32 x 20L per pallet) 5L (4 x 5L per shipper)

Active Constituents:

Formulation:

336g/L Mecoprop

Soluble Liquid (SL)

80g/L 2,4-D

40g/L Dicamba



At a Glance



Product Type:

Resistance group:

Pack Size:

Herbicide

Group 4 Herbicide

10L (75 x 10L per pallet)

Active Constituents:

Formulation:

336g/L Mecoprop,

80g/L MCPA

40g/L Dicamba

Soluble Liquid (SL)

Complete Broadleaf Weed Control

Methar Tri-Kombi and Sportsground is registered for use on the following weeds at the rate of 4.2L/ha

Can be used at 3L/ha for newely sown turf (refer ot label)

Clovers	Chickweed	Bindii	Cobblers Peg
Creeping Oxalis	Lambs Tongue	Fumitory	Capeweed
Pearlwort	Catsear	Staggerweeds	Wireweed
Mouse-eared chickweed	Fat Hen	Mallow	











Best Use Practices

- 1. Ensure weeds are actively growing before herbicide application and turf is not mowed 2 days prior or after to ensure best possible results. This allows better absorption.
- 2. Allow at least 4-10 hours before irrigation or rain to ensure that the spray will be absorbed, the longer the better.
- 3. Avoid spraying in very hot dry conditions and if frost is likley to occur.
- 4. Using a suitable surfactant (such as Filmstar) on difficult weeds (must be used on oxalis). This will increase the distribution and adhesion of the spray hence increasing the efficacy.
- 5. The timing of sprays can be crucial. Weeds are always best sprayed when they are young; at this stage they are most susceptible to herbicides.
- 6. Spray until weeds and stem are just wet. Spraying beyond this point will encourage run off away from the target area with loss of efficacy.
- 7. Seeding can take place the day after. However do not apply to newly seeded turf for at least 4 weeks after first mowing (usually 8-12 weeks after germination). Then use the low rate of 30mL/100m².


FilmStar - ideal to add to our herbicide products for better sticking and spreading

Methar Tri-Kombi & Sportsground are available from









"Good people, good products and good service with a dedication to the turf industry." - Gary Dempsey (Retired Superintendent NSW Golf Club)